


NASA/TM-2004-213007



Contributions of Spherical Harmonics to Magnetic and Gravitational Fields

Carlos M. Roithmayr
Langley Research Center, Hampton, Virginia

March 2004

The NASA STI Program Office ... in Profile

Since its founding, NASA has been dedicated to the advancement of aeronautics and space science. The NASA Scientific and Technical Information (STI) Program Office plays a key part in helping NASA maintain this important role.

The NASA STI Program Office is operated by Langley Research Center, the lead center for NASA's scientific and technical information. The NASA STI Program Office provides access to the NASA STI Database, the largest collection of aeronautical and space science STI in the world. The Program Office is also NASA's institutional mechanism for disseminating the results of its research and development activities. These results are published by NASA in the NASA STI Report Series, which includes the following report types:


- **TECHNICAL PUBLICATION.** Reports of completed research or a major significant phase of research that present the results of NASA programs and include extensive data or theoretical analysis. Includes compilations of significant scientific and technical data and information deemed to be of continuing reference value. NASA counterpart of peer-reviewed formal professional papers, but having less stringent limitations on manuscript length and extent of graphic presentations.
- **TECHNICAL MEMORANDUM.** Scientific and technical findings that are preliminary or of specialized interest, e.g., quick release reports, working papers, and bibliographies that contain minimal annotation. Does not contain extensive analysis.
- **CONTRACTOR REPORT.** Scientific and technical findings by NASA-sponsored contractors and grantees.
- **CONFERENCE PUBLICATION.** Collected papers from scientific and technical conferences, symposia, seminars, or other meetings sponsored or co-sponsored by NASA.
- **SPECIAL PUBLICATION.** Scientific, technical, or historical information from NASA programs, projects, and missions, often concerned with subjects having substantial public interest.
- **TECHNICAL TRANSLATION.** English-language translations of foreign scientific and technical material pertinent to NASA's mission.

Specialized services that complement the STI Program Office's diverse offerings include creating custom thesauri, building customized databases, organizing and publishing research results ... even providing videos.

For more information about the NASA STI Program Office, see the following:

- Access the NASA STI Program Home Page at <http://www.sti.nasa.gov>
- E-mail your question via the Internet to help@sti.nasa.gov
- Fax your question to the NASA STI Help Desk at (301) 621-0134
- Phone the NASA STI Help Desk at (301) 621-0390
- Write to:
NASA STI Help Desk
NASA Center for AeroSpace Information
7121 Standard Drive
Hanover, MD 21076-1320

NASA/TM-2004-213007



Contributions of Spherical Harmonics to Magnetic and Gravitational Fields

Carlos M. Roithmayr
Langley Research Center, Hampton, Virginia

National Aeronautics and
Space Administration

Langley Research Center
Hampton, Virginia 23681-2199

March 2004

Acknowledgment

The material presented here was written on December 4, 1992; it was subsequently distributed as a Johnson Space Center internal memorandum numbered EG-96-02 and dated January 23, 1996.

Available from:

NASA Center for AeroSpace Information (CASI)
7121 Standard Drive
Hanover, MD 21076-1320
(301) 621-0390

National Technical Information Service (NTIS)
5285 Port Royal Road
Springfield, VA 22161-2171
(703) 605-6000

Contributions of Spherical Harmonics to Magnetic and Gravitational Fields

Carlos Roithmayr

Introduction

Gravitational forces are of cardinal importance in the dynamics of spacecraft; magnetic attractions sometime play a significant role also, as was the case with the Long Duration Exposure Facility, and as is now true for the first segment of Space Station Freedom. Both satellites depend on gravitational moment and a device known as a magnetic damper to stabilize their orientation.

Magnetic fields are mathematically similar to gravitational fields in one important respect: each can be regarded as a gradient of a potential function that, in turn, can be described as an infinite series of spherical harmonics. Consequently, the two fields can be computed, in part, with quantities that need only be evaluated once, resulting in a savings of time when both fields are needed.

The objective of this material is to present magnetic field and gravitational force expressions, and point out the terms that belong to both—this is accomplished in Sections 1 and 2. Section 3 contains the deductive reasoning with which one obtains the expressions of interest. Finally, examples in Section 4 show these equations can be used to reproduce others that arise in connection with special cases such as the magnetic field produced by a tilted dipole, and gravitational force exerted by an oblate spheroid. The mathematics are discussed in the context of terrestrial fields; however, by substituting appropriate constants, the results can be made applicable to fields belonging to other celestial bodies.

The expressions presented here share the characteristics of algorithms set forth in Refs. [1]–[4] for computing gravitational force. In particular, computation is performed speedily by means of recursion formulae, and the expressions do not suffer from the shortcoming of a singularity when evaluated at points that lie on the polar axis.

1 Contributions to Magnetic Field

The magnetic field vector \mathbf{B} at a point Q above the Earth's surface can be expressed as the sum

$$\mathbf{B} = \sum_{n=1}^{\infty} \sum_{m=0}^n \mathbf{B}_{n,m} \quad (1)$$

where $\mathbf{B}_{n,m}$ represents the contribution to \mathbf{B} of the spherical harmonic of degree n and order m , and is given by

$$\mathbf{B}_{n,m} = \frac{K_{n,m} a^{n+2}}{R^{n+m+1}} \left\{ \frac{g_{n,m} \mathcal{C}_m + h_{n,m} \mathcal{S}_m}{R} [(s_\lambda A_{n,m+1} + (n+m+1)A_{n,m}) \hat{\mathbf{r}} - A_{n,m+1} \hat{\mathbf{e}}_3] \right. \\ \left. - mA_{n,m} [(g_{n,m} \mathcal{C}_{m-1} + h_{n,m} \mathcal{S}_{m-1}) \hat{\mathbf{e}}_1 + (h_{n,m} \mathcal{C}_{m-1} - g_{n,m} \mathcal{S}_{m-1}) \hat{\mathbf{e}}_2] \right\} \quad (2)$$

Here, a is the mean radius of the Earth (6371 km: p. 25, Ref. [5]) adopted for the International Geomagnetic Reference Field; R is the magnitude of \mathbf{R} , the position vector from the geometric center of the Earth to Q ; $\hat{\mathbf{r}}$ is a unit vector in the direction of \mathbf{R} ; and $g_{n,m}$ and $h_{n,m}$ represent Gauss coefficients of degree n and order m . Mutually perpendicular unit vectors $\hat{\mathbf{e}}_1$, $\hat{\mathbf{e}}_2$, and $\hat{\mathbf{e}}_3$ are fixed in the Earth: $\hat{\mathbf{e}}_1$ lies in the equatorial plane parallel to a line intersecting Earth's geometric center and the Greenwich meridian, $\hat{\mathbf{e}}_3$ is in the direction of the north polar axis, and $\hat{\mathbf{e}}_2 = \hat{\mathbf{e}}_3 \times \hat{\mathbf{e}}_1$.

The coefficients $K_{n,m}$ in Eq. (2) are given by

$$K_{n,0} = 1 \quad (n = 1, \dots, \infty) \quad (3)$$

whenever $m = 0$ (in addition, $K_{1,1} = 1$); otherwise, they can be obtained with one of two recursion formulae:

$$K_{n,m} = \left(\frac{n-m}{n+m} \right)^{\frac{1}{2}} K_{n-1,m} \quad [m = 1, \dots, \infty; n \geq (m+1)] \quad (4)$$

$$K_{n,m} = [(n+m)(n-m+1)]^{-\frac{1}{2}} K_{n,m-1} \quad (m = 2, \dots, \infty; n \geq m) \quad (5)$$

$A_{n,m}$ and $A_{n,m+1}$ are referred to as derived Legendre polynomials of degree n , and orders m and $m+1$ respectively. The notation $A_{n,m}(u)$ is used to indicate that $A_{n,m}$ is, in general, a function of an argument u ; however, for the sake of clarity, the arguments of $A_{n,m}$ and $A_{n,m+1}$ do not appear explicitly in Eq. (2). When $n = m$, the derived Legendre polynomial is given by a simple expression that can be written in recursive form.

$$A_{n,n} = (1)(3)(5) \cdots (2n-1) \quad (n = 1, \dots, \infty) \\ = (2n-1)A_{n-1,n-1} \quad (n = 2, \dots, \infty) \quad (6)$$

It should be noted that $A_{1,1}$, $A_{2,2}$, \dots are not functions of an argument; therefore, they need only be computed once to provide the seeds for a recursion relation shown in Ref. [6]

to give numerically stable and precise results:

$$A_{n,m}(u) = \frac{1}{n-m} [(2n-1)uA_{n-1,m}(u) - (n+m-1)A_{n-2,m}(u)]$$

$$[m = 0, \dots, \infty; n \geq (m+1)] \quad (7)$$

$A_{n,m}(u)$ vanishes when $m > n$, and $A_{0,0}(u) = 1$. Arguments of the derived Legendre polynomials in Eq. (2) are $\sin \lambda$ (shortened to s_λ), where λ is the geographic latitude of Q . $s_\lambda = \hat{\mathbf{r}} \cdot \hat{\mathbf{e}}_3$.

The symbols \mathcal{S}_m and \mathcal{C}_m appearing in Eq. (2) will be defined presently. Their values for $m = 0$ and $m = 1$,

$$\mathcal{S}_0 = 0, \quad \mathcal{C}_0 = 1 \quad (8)$$

$$\mathcal{S}_1 = \mathbf{R} \cdot \hat{\mathbf{e}}_2, \quad \mathcal{C}_1 = \mathbf{R} \cdot \hat{\mathbf{e}}_1 \quad (9)$$

provide a means of starting the recursion relations given on p. 4 of Ref. [2].

$$\mathcal{S}_m = \mathcal{S}_1 \mathcal{C}_{m-1} + \mathcal{C}_1 \mathcal{S}_{m-1}, \quad \mathcal{C}_m = \mathcal{C}_1 \mathcal{C}_{m-1} - \mathcal{S}_1 \mathcal{S}_{m-1} \quad (10)$$

2 Contributions to Gravitational Field

Like the magnetic field vector, the gravitational force per unit mass \mathbf{f} exerted by the Earth at a point Q above the Earth's surface, can be regarded as a sum:

$$\mathbf{f} = -\frac{\mu}{R^2} \hat{\mathbf{r}} + \sum_{n=2}^{\infty} \sum_{m=0}^n \mathbf{f}_{n,m} \quad (11)$$

where μ , the gravitational parameter, is the product of G , the universal gravitational constant, and M , the Earth's mass; the position vector from Earth's mass center to Q has magnitude R and the direction of unit vector $\hat{\mathbf{r}}$; and $\mathbf{f}_{n,m}$ represents the contribution to \mathbf{f} of the spherical harmonic of degree n and order m , given by

$$\mathbf{f}_{n,m} = \frac{\mu R_\oplus^n}{R^{n+m+1}} \left\{ \frac{C_{n,m} \mathcal{C}_m + S_{n,m} \mathcal{S}_m}{R} [A_{n,m+1} \hat{\mathbf{e}}_3 - (s_\lambda A_{n,m+1} + (n+m+1) A_{n,m}) \hat{\mathbf{r}}] \right.$$

$$\left. + mA_{n,m} [(C_{n,m} \mathcal{C}_{m-1} + S_{n,m} \mathcal{S}_{m-1}) \hat{\mathbf{e}}_1 + (S_{n,m} \mathcal{C}_{m-1} - C_{n,m} \mathcal{S}_{m-1}) \hat{\mathbf{e}}_2] \right\} \quad (12)$$

R_\oplus is the mean equatorial radius of the Earth (6,378,139 m), and $C_{n,m}$ and $S_{n,m}$ are gravitational coefficients of degree n and order m . Mutually perpendicular unit vectors $\hat{\mathbf{e}}_1$, $\hat{\mathbf{e}}_2$, and $\hat{\mathbf{e}}_3$ have the same directions as in Eq. (2). $A_{n,m}$, \mathcal{S}_m , and \mathcal{C}_m have the same meanings as in Eq. (2); consequently, they need only be computed once when constructing magnetic and gravitational fields.

It is not too difficult to show that Eq. (11) and Eq. (12), which has the form $\mathbf{f}_{n,m} = x_{n,m} \hat{\mathbf{e}}_1 + y_{n,m} \hat{\mathbf{e}}_2 + z_{n,m} \hat{\mathbf{e}}_3 + w_{n,m} \hat{\mathbf{r}}$ [Eq. (2) has this form as well], are in concert with Eq. (31) of Ref. [1], whose form is $\mathbf{f} = X \hat{\mathbf{e}}_1 + Y \hat{\mathbf{e}}_2 + Z \hat{\mathbf{e}}_3 + W \hat{\mathbf{r}}$: in other words, $W = -\mu/R^2 + \sum_{n=2}^{\infty} \sum_{m=0}^n w_{n,m}$, $X = \sum_{n=2}^{\infty} \sum_{m=0}^n x_{n,m}$, and so forth. (The sum to the right of the second = sign in Eqs. (30a) of Ref. [1] should be preceded by a minus sign.)

3 Derivations

Eq. (2) is similar in form to Eq. (12) because \mathbf{B} and \mathbf{f} are each obtained by differentiating a potential function that is expressed in terms of spherical harmonics. A potential function associated with the magnetic field is given below, followed by steps in the differentiation process that results in Eq. (2). Next, a gravitational potential function is displayed. The differentiation process that leads to Eq. (12) is virtually identical to that used with the magnetic potential; therefore, it does not bear repeating.

3.1 Magnetic Field

The magnetic field vector \mathbf{B} is related to V_β , a scalar geomagnetic potential, by the equation

$$\mathbf{B} = -\nabla V_\beta \quad (13)$$

where ∇ denotes differentiation with respect to \mathbf{R} : \mathbf{B} is the negative of the spatial gradient of V_β . Section 2.9 of Ref. [7] contains a thorough explanation of how one differentiates with respect to a vector. The potential V_β satisfies Laplace's equation and, consequently, is expressed in Eq. (2.13) of Ref. [5], and in Eq. (9.0,6) of Ref. [8], as an infinite series of spherical harmonics.

$$V_\beta = a \sum_{n=1}^{\infty} \left(\frac{a}{R}\right)^{n+1} \sum_{m=0}^n P_n^m(\cos \theta) \left[g_{n,m} \cos m\phi + h_{n,m} \sin m\phi \right] \quad (14)$$

where a , R , $g_{n,m}$, and $h_{n,m}$ have the same meanings as in Eq. (2), and ϕ is the geographic longitude of Q measured Eastward from the Greenwich meridian. $P_n^m(\cos \theta)$ are Schmidt functions (they will be discussed directly) whose arguments are $\cos \theta$, where θ is the geographic co-latitude of Q . Now, Eq. (14) can be rewritten as

$$V_\beta = \sum_{n=1}^{\infty} \sum_{m=0}^n V_{n,m} \quad (15)$$

so long as $V_{n,m}$ is defined as

$$V_{n,m} \triangleq a \left(\frac{a}{R}\right)^{n+1} P_n^m(\cos \theta) \left[g_{n,m} \cos m\phi + h_{n,m} \sin m\phi \right] \quad (16)$$

For the sake of clarity, we dispense with the subscript β for $V_{n,m}$. Thus, in view of Eqs. (13) and (15), \mathbf{B} can be expressed as in Eq. (1)

$$\mathbf{B} = -\nabla V_\beta = -\nabla \sum_{n=1}^{\infty} \sum_{m=0}^n V_{n,m} = \sum_{n=1}^{\infty} \sum_{m=0}^n \mathbf{B}_{n,m} \quad (17)$$

with the contribution to \mathbf{B} of the spherical harmonic of degree n and order m defined simply as

$$\mathbf{B}_{n,m} \triangleq -\nabla V_{n,m} \quad (18)$$

The Schmidt functions P_n^m are of degree n and order m , and are related [Ref. [9], Chapter 9, Eq. (13)] to $P_{n,m}$, associated Legendre functions of the first kind.

$$P_n^m(\cos \theta) = \begin{cases} \left[2 \frac{(n-m)!}{(n+m)!}\right]^{\frac{1}{2}} P_{n,m}(\cos \theta) & (m = 1, \dots, n) \\ P_{n,m}(\cos \theta) & (m = 0) \end{cases} \quad (19)$$

Angle θ , the co-latitude of Q , is the complement of λ , the latitude of Q ; therefore, the argument $\cos \theta$ of P_n^m and $P_{n,m}$ can be replaced with s_λ , which is shorthand for $\sin \lambda$. Products of $P_{n,m}(s_\lambda)$ with $\cos m\phi$, or with $\sin m\phi$, are referred to as tesseral harmonics of degree n and order m : they are functions of latitude and longitude. Tesseral harmonics of order zero are called zonal harmonics, and are independent of longitude.

In the interest of economy in written symbols, the coefficients of $P_{n,m}$ in Eqs. (19) can be labeled $K_{n,m}$ [see Eq. (2)] and defined as

$$K_{n,m} \triangleq \begin{cases} \left[2 \frac{(n-m)!}{(n+m)!}\right]^{\frac{1}{2}} & (m = 1, \dots, n) \\ 1 & (m = 0) \end{cases} \quad (20)$$

The recursive relationships in Eqs. (4) and (5) follow from the first of these definitions; Eqs. (3) from the second.

The associated Legendre functions $P_{n,m}$ are, in turn, related to derived Legendre polynomials $A_{n,m}$ by Eq. (9b) of Ref. [6] (see also p. 3 of Ref. [2]).

$$P_{n,m}(s_\lambda) = (\cos \lambda)^m A_{n,m}(s_\lambda) = \left(\frac{\rho}{R}\right)^m A_{n,m}(s_\lambda) \quad (21)$$

where ρ is related to \mathbf{R} by the definition

$$\rho^2 \triangleq (\mathbf{R} \cdot \hat{\mathbf{e}}_1)^2 + (\mathbf{R} \cdot \hat{\mathbf{e}}_2)^2 \quad (22)$$

In words, ρ is the magnitude of the projection of \mathbf{R} onto Earth's equatorial plane.

The symbols \mathcal{S}_m and \mathcal{C}_m appearing in Eq. (2) are defined on p. 4 of Ref. [2].

$$\mathcal{S}_m \triangleq \rho^m \sin m\phi, \quad \mathcal{C}_m \triangleq \rho^m \cos m\phi \quad (23)$$

where ϕ has the same meaning as in Eq. (14).

Differentiation of $V_{n,m}$ with respect to \mathbf{R} is facilitated by substitution from Eqs. (19), (20), (21), and (23) into (16).

$$\begin{aligned} V_{n,m} &= a \left(\frac{a}{R}\right)^{n+1} K_{n,m} \left(\frac{\rho}{R}\right)^m A_{n,m}(s_\lambda) [g_{n,m} \cos m\phi + h_{n,m} \sin m\phi] \\ &= K_{n,m} a^{n+2} A_{n,m} \frac{g_{n,m} \mathcal{C}_m + h_{n,m} \mathcal{S}_m}{R^{n+m+1}} \end{aligned} \quad (24)$$

$K_{n,m}$, a , $g_{n,m}$, and $h_{n,m}$ are independent of \mathbf{R} , so obtaining an expression for the gradient $\nabla V_{n,m}$ becomes a matter of producing the four gradients $\nabla R^{-(n+m+1)}$, $\nabla A_{n,m}$, $\nabla \mathcal{C}_m$, and $\nabla \mathcal{S}_m$.

Eqs. (2.9.7)–(2.9.9) of Ref. [7] provide the machinery with which all the required gradients can be formed:

$$\nabla \mathbf{R} = \underline{\mathbf{U}} \quad (25)$$

$$\nabla R = \hat{\mathbf{r}} \quad (26)$$

$$\nabla \hat{\mathbf{r}} = R^{-1}(\underline{\mathbf{U}} - \hat{\mathbf{r}}\hat{\mathbf{r}}) \quad (27)$$

where $\underline{\mathbf{U}}$ is the unit dyadic. (When $\underline{\mathbf{U}}$ is pre or post multiplied by any vector \mathbf{v} , the result is \mathbf{v} , that is, $\underline{\mathbf{U}} \cdot \mathbf{v} = \mathbf{v} \cdot \underline{\mathbf{U}} = \mathbf{v}$.) The first gradient is fashioned easily with the help of Eq. (26).

$$\nabla R^{-(n+m+1)} = -(n+m+1)R^{-(n+m+2)}\hat{\mathbf{r}} \quad (28)$$

Referring to Eq. (10) of Ref. [6], one can write $\nabla A_{n,m}(s_\lambda) = A_{n,m+1}\nabla s_\lambda$, express s_λ in terms of $\hat{\mathbf{r}}$ and $\hat{\mathbf{e}}_3$ ($\hat{\mathbf{e}}_3$ is independent of \mathbf{R}) in order to expedite differentiation with respect to \mathbf{R} by means of Eq. (27),

$$\nabla s_\lambda = \nabla(\hat{\mathbf{r}} \cdot \hat{\mathbf{e}}_3) = R^{-1}(\underline{\mathbf{U}} - \hat{\mathbf{r}}\hat{\mathbf{r}}) \cdot \hat{\mathbf{e}}_3 = [\hat{\mathbf{e}}_3 - (\hat{\mathbf{r}} \cdot \hat{\mathbf{e}}_3)\hat{\mathbf{r}}]/R = (\hat{\mathbf{e}}_3 - s_\lambda\hat{\mathbf{r}})/R \quad (29)$$

and obtain the second gradient.

$$\nabla A_{n,m}(s_\lambda) = (\hat{\mathbf{e}}_3 - s_\lambda\hat{\mathbf{r}})A_{n,m+1}/R \quad (30)$$

Finally, the remaining two gradients can be produced from Eqs. (23) with the aid of Eqs. (22) and (25),

$$\nabla \mathcal{S}_m = m(\mathcal{S}_{m-1}\hat{\mathbf{e}}_1 + \mathcal{C}_{m-1}\hat{\mathbf{e}}_2), \quad \nabla \mathcal{C}_m = m(\mathcal{C}_{m-1}\hat{\mathbf{e}}_1 - \mathcal{S}_{m-1}\hat{\mathbf{e}}_2) \quad (31)$$

as is suggested by Eqs. (7) in Ref. [2].

The foregoing material puts one into position to express $\mathbf{B}_{n,m}$ as in Eq. (2) by substituting from Eq. (24) into (18), and then employing the right hand members of Eqs. (28), (30), and (31).

3.2 Gravitational Field

The force \mathbf{f} in a gravitational field is analogous to the vector \mathbf{B} in a magnetic field: both are spatial gradients of scalar potential functions. At a point Q above Earth's surface, a particle of unit mass is subject to a gravitational force \mathbf{f} given by

$$\mathbf{f} = \nabla V_\gamma \quad (32)$$

where V_γ is a scalar gravitational potential, and ∇ denotes differentiation with respect to \mathbf{R} , the position vector from Earth's mass center to Q . Like V_β , V_γ satisfies Laplace's equation and can be expressed as an infinite sum of spherical harmonics. Eq. (2.13.12) of

Ref. [7] is one such example, written in the form adopted as a standard by the International Astronomical Union:

$$\begin{aligned} V_\gamma &= \frac{\mu}{R} \left\{ 1 + \sum_{n=2}^{\infty} \left(\frac{R_\oplus}{R} \right)^n \sum_{m=0}^n P_{n,m}(\sin \lambda) [C_{n,m} \cos m\phi + S_{n,m} \sin m\phi] \right\} \\ &= \frac{\mu}{R} + \sum_{n=2}^{\infty} \sum_{m=0}^n V_{n,m} \end{aligned} \quad (33)$$

where μ , R_\oplus , $C_{n,m}$, and $S_{n,m}$ have the same meanings as in Eq. (12), and Eqs. (21) and (23) can be engaged to define $V_{n,m}$:

$$V_{n,m} \triangleq \mu R_\oplus^n A_{n,m} \frac{C_{n,m} \mathcal{C}_m + S_{n,m} \mathcal{S}_m}{R^{n+m+1}} \quad (34)$$

Thus, Eq. (11) is the result of substituting from Eq. (33) into (32).

$$\mathbf{f} = \nabla \frac{\mu}{R} + \sum_{n=2}^{\infty} \sum_{m=0}^n \nabla V_{n,m} = -\frac{\mu}{R^2} \hat{\mathbf{r}} + \sum_{n=2}^{\infty} \sum_{m=0}^n \mathbf{f}_{n,m} \quad (35)$$

and the expression for $\mathbf{f}_{n,m}$ in Eq. (12) follows from differentiation of the right hand member of Eq. (34) [which is of the same form as Eq. (24)] in the much the same way as $\mathbf{B}_{n,m}$ results from differentiation of the right side of Eq. (24). μ , R_\oplus , $C_{n,m}$, and $S_{n,m}$ are all independent of \mathbf{R} .

4 Examples

Although Eqs. (2) and (12) are intended to be evaluated numerically, rather than analytically, they can be used to produce recognizable expressions in a few important special cases. In this Section, Eqs. (2) and (12) are exercised through applications to a magnetic field associated with a tilted dipole, to the gravitational force exerted by a planet whose mass is distributed symmetrically about its polar axis, and to the gravitational attraction of an oblate spheroid—all of which have been dealt with elsewhere.

4.1 Magnetic Field of a Tilted Dipole

The geomagnetic field can be modeled as a tilted dipole by adding the contributions of spherical harmonics of degree 1, and orders 0 and 1. According to Ref. [5], approximately 90% of the magnetic field at the Earth's surface is described by this model. The derived Legendre polynomials identified with the dipole are [see Eqs. (6) and (7)] simply

$$A_{1,0}(s_\lambda) = s_\lambda, \quad A_{1,1}(s_\lambda) = 1, \quad A_{1,2}(s_\lambda) = 0 \quad (36)$$

and Eq. (20) yields the coefficients

$$K_{1,0} = K_{1,1} = 1 \quad (37)$$

Substitution from Eqs. (36), (37), (8), and (9) into (2) leads, for $n = 1$, $m = 0$, and $m = 1$, to

$$\mathbf{B}_{1,0} = \left(\frac{a}{R}\right)^3 g_{1,0}(3s_\lambda \hat{\mathbf{r}} - \hat{\mathbf{e}}_3) \quad (38)$$

$$\mathbf{B}_{1,1} = \left(\frac{a}{R}\right)^3 [3(g_{1,1} \hat{\mathbf{r}} \cdot \hat{\mathbf{e}}_1 + h_{1,1} \hat{\mathbf{r}} \cdot \hat{\mathbf{e}}_2) \hat{\mathbf{r}} - (g_{1,1} \hat{\mathbf{e}}_1 + h_{1,1} \hat{\mathbf{e}}_2)] \quad (39)$$

and then, after recognizing that $s_\lambda = \hat{\mathbf{r}} \cdot \hat{\mathbf{e}}_3$, to

$$\mathbf{B}_{1,0} + \mathbf{B}_{1,1} = \left(\frac{a}{R}\right)^3 [3(\hat{\mathbf{r}} \cdot \mathbf{M}) \hat{\mathbf{r}} - \mathbf{M}] \quad (40)$$

which is in agreement with Eq. (H-22) of Ref. [10] when \mathbf{M} , the terrestrial dipole moment, is defined as $\mathbf{M} \triangleq g_{1,1} \hat{\mathbf{e}}_1 + h_{1,1} \hat{\mathbf{e}}_2 + g_{1,0} \hat{\mathbf{e}}_3$.

4.2 Gravitational Force Exerted by an Axisymmetric Body

An expression for the gravitational force per unit mass, exerted by an axisymmetric body, is given in Ref. [11], and can be reproduced from Eqs. (11) and (12). The tesseral harmonics vanish from the gravitational potential as a consequence of axial symmetry, leaving only the zonal harmonics. Working only with terms of order 0, and substituting from Eq. (8) into (12), we get

$$\mathbf{f} = -\frac{\mu}{R^2} \hat{\mathbf{r}} + \sum_{n=2}^{\infty} \mathbf{f}_{n,0} = -\frac{\mu}{R^2} \hat{\mathbf{r}} - \mu \sum_{n=2}^{\infty} J_n \frac{R_\oplus^n}{R^{n+2}} \{A_{n,1} \hat{\mathbf{e}}_3 - [s_\lambda A_{n,1} + (n+1)A_{n,0}] \hat{\mathbf{r}}\} \quad (41)$$

where the zonal harmonic coefficients, $C_{n,0}$, have been renamed: $C_{n,0} \triangleq -J_n$. The derived Legendre polynomials are defined in Eq. (14) of Ref. [1] as $A_{n,m}(u) \triangleq d^m P_n(u)/du^m$, where $P_n(u)$ is a Legendre polynomial of degree n , with argument u . Denoting the first derivative (with respect to u) of $P_n(u)$ as $P'_n(u)$ leads to

$$\mathbf{f} = -\frac{\mu}{R^2} \hat{\mathbf{r}} - \mu \sum_{n=2}^{\infty} J_n \frac{R_\oplus^n}{R^{n+2}} \{P'_n \hat{\mathbf{e}}_3 - [s_\lambda P'_n + (n+1)P_n] \hat{\mathbf{r}}\} \quad (42)$$

which, upon use of the recursion relation $P'_{n+1}(u) = uP'_n(u) + (n+1)P_n(u)$, becomes the expression given on p. 407 of Ref. [11]:

$$\mathbf{f} = -\frac{\mu}{R^2} \left[\hat{\mathbf{r}} + \sum_{n=2}^{\infty} J_n \left(\frac{R_\oplus}{R}\right)^n (P'_n \hat{\mathbf{e}}_3 - P'_{n+1} \hat{\mathbf{r}}) \right] \quad (43)$$

4.3 Gravitational Force Exerted by an Oblate Spheroid

An oblate spheroid is an example of an axisymmetric body, and is often used to model the Earth. A celestial body's oblateness is represented by a zonal harmonic of degree 2, and is responsible for precessions in a satellite's orbit plane and argument of perigee. The derived Legendre polynomials identified with oblateness are [see Eqs. (6) and (7)] just

$$A_{2,0}(s_\lambda) = \frac{1}{2}(3s_\lambda^2 - 1), \quad A_{2,1}(s_\lambda) = 3s_\lambda \quad (44)$$

Substitution from Eqs. (44) and (8) into Eq. (12) yields, for $n = 2$ and $m = 0$,

$$\mathbf{f}_{2,0} = -\mu J_2 \frac{R_\oplus^2}{R^4} \left[3s_\lambda \hat{\mathbf{e}}_3 + \frac{3}{2}(1 - 5s_\lambda^2) \hat{\mathbf{r}} \right] \quad (45)$$

This expression for the contribution of oblateness to the gravitational force per unit mass, exerted by Earth, is in agreement with the vector equation (8.28) in Ref. [12] and, after a little manipulation, with the scalar equations for a_x , a_y , and a_z given on p. 3-1 of Ref. [3]. [$a_x = (-\mu \hat{\mathbf{r}}/R^2 + \mathbf{f}_{2,0}) \cdot \hat{\mathbf{e}}_1$, $a_y = (-\mu \hat{\mathbf{r}}/R^2 + \mathbf{f}_{2,0}) \cdot \hat{\mathbf{e}}_2$, and $a_z = (-\mu \hat{\mathbf{r}}/R^2 + \mathbf{f}_{2,0}) \cdot \hat{\mathbf{e}}_3$.]

References

- [1] Pines, S., "Uniform Representation of the Gravitational Potential and its Derivatives", *AIAA Journal*, Vol. 11, Nov. 1973, pp. 1508–1511.
- [2] Mueller, A. C., A Fast Recursive Algorithm For Calculating The Forces Due To The Geopotential, NASA Johnson Space Center Internal Note No. 75-FM-42, June 9, 1975.
- [3] Lear, W. M., The Gravitational Acceleration Equations, NASA Johnson Space Center, 86-FM-15, April, 1986.
- [4] Gottlieb, R. G., A Fast Recursive Singularity Free Algorithm for Calculating the First and Second Derivatives of the Geopotential, McDonnell Douglas Astronautics Co., Houston, Tx., 3.4-WP-FM8-EA-01, May, 1988.
- [5] Merrill, R. T., and McElhinny, M. W., *The Earth's Magnetic Field: Its History, Origin and Planetary Perspective*, Academic Press, New York, 1983.
- [6] Lundberg, J. B., and Schutz, B. E., "Recursion Formulas of Legendre Functions for Use with Nonsingular Geopotential Models", *Journal of Guidance, Control, and Dynamics*, Vol. 11, Jan–Feb 1988, pp. 32–38.
- [7] Kane, T. R., Likins, P. W., and Levinson, D. A., *Spacecraft Dynamics*, McGraw-Hill, New York, 1983.
- [8] *Space Station Program Natural Environment Definition for Design*, SSP 30425 Rev. A, NASA Space Station Freedom Program Office, Reston, Va., June 1991.

- [9] Haymes, R. C., *Introduction to Space Science*, Wiley, New York, 1971.
- [10] Plett, M., Magnetic Field Models, Appendix H in Wertz, J. R. (ed.), *Spacecraft Attitude Determination and Control*, D. Reidel Publishing Co., Dordrecht, Holland, 1978.
- [11] Battin, R. H., *An Introduction to the Mathematics and Methods of Astrodynamics*, AIAA, New York, 1987.
- [12] Kaplan, M. H., *Modern Spacecraft Dynamics and Control*, John Wiley & Sons, New York, 1976.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE (DD-MM-YYYY) 01-03-2004		2. REPORT TYPE Technical Memorandum		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Contributions of Spherical Harmonics to Magnetic and Gravitational Fields			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) Carlos M. Roithmayr			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER 23-319-30-02		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) NASA Langley Research Center Hampton, VA 23681-2199			8. PERFORMING ORGANIZATION REPORT NUMBER L-18358		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Washington, DC 20546-0001			10. SPONSOR/MONITOR'S ACRONYM(S) NASA		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S) NASA/TM-2004-213007		
12. DISTRIBUTION/AVAILABILITY STATEMENT Unclassified-Unlimited Subject Category 90 Availability: NASA CASI (301) 621-0390 Distribution: Nonstandard					
13. SUPPLEMENTARY NOTES An electronic version can be found at http://techreports.larc.nasa.gov/trs/ or http://ntrs.nasa.gov .					
14. ABSTRACT Analysis of the behavior of a spacecraft often requires a model of the gravitational field in which it operates, or the magnetic field, or both. Each type of vector field can be regarded as the gradient of a scalar potential function expressed as an infinite series of spherical harmonics. The contribution from the harmonic of degree n and order m to each field is worked out, and the terms common to both fields are identified. Expressions for each contribution are evaluated speedily by means of recursion formulae and by calculating the common terms just once; the expressions are free of singularity at points on the polar axis of the body that is the source of the field. Examples show the expressions can be used to reproduce existing results in special cases such as the magnetic field of a tilted dipole, or the gravitational field exerted by an oblate spheroid.					
15. SUBJECT TERMS Spherical harmonics, tesseral harmonics, zonal harmonics, gravitational field, magnetic field, derived Legendre polynomial, associated Legendre functions of the first kind, recursion formulae, potential function, Schmidt function, spatial gradient, gravitational coefficients, Gauss coefficients, International Geomagnetic Reference Field, tilted dipole, oblate spheroid.					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 15	19a. NAME OF RESPONSIBLE PERSON STI Help Desk (email: help@sti.nasa.gov)
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			